


## pFusionRed-CDC42 vector

The vector sequence has been compiled using the information from sequence databases, published literature, and other sources, together with partial sequences obtained by Evrogen. This vector has not been completely sequenced.



For vector sequence, please visit our Web site at <http://www.evrogen.com/products/vectors.shtml>

### Location of features

P<sub>CMV IE</sub>: 1-589  
 Enhancer region: 59-465  
 TATA box: 554-560  
 Transcription start point: 583  
 Kozak consensus translation initiation site: 606-616  
 FusionRed  
 Start codon (ATG): 613-615  
 Last amino acid in FusionRed: 1306-1308  
 Human CDC42: 1351-1926  
 Stop codon: 1924-1926  
 SV40 early mRNA polyadenylation signal  
 Polyadenylation signals: 2087-2092 & 2116-2121  
 mRNA 3' ends: 2125 & 2137  
 f1 single-strand DNA origin: 2184-2639  
 Bacterial promoter for expression of Kan<sup>r</sup> gene  
 -35 region: 2701-2706; -10 region: 2724-2729  
 Transcription start point: 2736  
 SV40 origin of replication: 2980-3115  
 SV40 early promoter  
 Enhancer (72-bp tandem repeats): 2813-2884 & 2885-2956  
 21-bp repeats: 2960-2980, 2981-3001 & 3003-3023  
 Early promoter element: 3036-3042  
 Major transcription start points: 3032, 3070, 3076 & 3081  
 Kanamycin/neomycin resistance gene  
 Neomycin phosphotransferase coding sequences:  
 Start codon (ATG): 3164-3166; Stop codon: 3956-3958  
 G->A mutation to remove Pst I site: 3346  
 C->A (Arg to Ser) mutation to remove BssH II site: 3692  
 Herpes simplex virus (HSV) thymidine kinase (TK) polyadenylation signal  
 Polyadenylation signals: 4194-4199 & 4207-4212  
 pUC plasmid replication origin: 4543-5186

Product	Cat.#	Size
pFusionRed-CDC42 vector	<b>FP435</b>	20 µg
Vector type	mammalian expression vector	
Reporter	FusionRed	
Reporter codon usage	mammalian	
Promoter for FusionRed	P <sub>CMV IE</sub>	
Host cells	mammalian	
Selection	prokaryotic - kanamycin eukaryotic - neomycin (G418)	
Replication	prokaryotic - pUC ori eukaryotic - SV40 ori	
Use	red fluorescent labeling of CDC42	

### Vector description

pFusionRed-CDC42 is a mammalian expression vector encoding FusionRed-CDC42 fusion protein. The vector can be used for fluorescent labeling of CDC42 in living cells.

FusionRed codon usage is optimized for high expression in mammalian cells (humanized) [Haas et al. 1996]. CDC42 is fused to the FusionRed C-terminus. To increase mRNA translation efficiency, Kozak consensus translation initiation site is generated upstream of the FusionRed-CDC42 coding sequence [Kozak 1987].

pFusionRed-CDC42 vector can be used as a source of FusionRed-CDC42 hybrid sequence. The vector backbone contains unique restriction sites that permit its excision and further insertion into expression vector of choice.

**Note:** The plasmid DNA was isolated from dam<sup>+</sup>-methylated *E. coli*. Therefore some restriction sites are blocked by methylation. If you wish to digest the vector using such sites you will need to transform the vector into a dam<sup>-</sup> host and make fresh DNA.

The vector backbone contains immediate early promoter of cytomegalovirus (P<sub>CMV IE</sub>) for protein expression, SV40 origin for replication in mammalian cells expressing SV40 T-antigen, pUC origin of replication for propagation in *E. coli*, and f1 origin for single-stranded DNA production. SV40 polyadenylation signals (SV40 poly A) direct proper processing of the 3'-end of the reporter mRNA.

SV40 early promoter (P<sub>SV40</sub>) provides neomycin resistance gene (Neo<sup>r</sup>) expression to select stably transfected eukaryotic cells using G418. Bacterial promoter (P) provides kanamycin resistance gene expression (Kan<sup>r</sup>) in *E. coli*. Kan<sup>r</sup>/Neo<sup>r</sup> gene is linked with herpes simplex virus (HSV) thymidine kinase (TK) polyadenylation signals.

### Expression in mammalian cells

pFusionRed-CDC42 vector can be transfected into mammalian cells by any known transfection method. CMV promoter provides strong, constitutive expression of the FusionRed-CDC42 fusion in eukaryotic cells. If required, stable transformants can be selected using G418 [Gorman 1985].

### Propagation in *E. coli*

Suitable host strains for propagation in *E. coli* include DH5alpha, HB101, XL1-Blue, and other general purpose strains. Plasmid incompatibility group is pMB1/ColE1. The vector confers resistance to kanamycin (30 µg/ml) to *E. coli* hosts. Copy number in *E. coli* is about 500.

### References

- Gorman, C. (1985). "High efficiency gene transfer into mammalian cells." In: *DNA cloning: A Practical Approach, Vol. II*. Ed. by Glover. (IRL Press, Oxford, U.K.) Pp. 143-190.
- Haas, J. et al. (1996) "Codon usage limitation in the expression of HIV-1 envelope glycoprotein." *Curr Biol*, 6 (3): 315-324 / pmid: 8805248
- Kozak, M. (1987) "An analysis of 5'-noncoding sequences from 699 vertebrate messenger RNAs." *Nucleic Acids Res*, 15 (20): 8125-8148 / pmid: 3313277

### Notice to Purchaser:

FusionRed-related materials (also referred to as "Products") are intended for research use only. The Products are covered by European Pat. 1994149 and other Evrogen Patents and/or Patent applications pending. By use of these Products, you accept the terms and conditions of the applicable Limited Use Label License #001: <http://www.evrogen.com/products/Evrogen-FP-license.shtml>.

The CMV promoter is covered under U.S. Patents 5,168,062 and 5,385,839, and its use is permitted for research purposes only. Any other use of the CMV promoter requires a license from the University of Iowa Research Foundation, 214 Technology Innovation Center, Iowa City, IA 52242.

**MSDS information** is available at <http://www.evrogen.com/MSDS.shtml>