


pFusionRed-ER vector

The vector sequence has been compiled using the information from sequence databases, published literature, and other sources, together with partial sequences obtained by Evrogen. This vector has not been completely sequenced.



For vector sequence, please visit our Web site at <http://www.evrogen.com/products/vectors.shtml>

Location of features

P_{CMV IE}: 1-589
 Enhancer region: 59-465
 TATA box: 554-560
 Transcription start point: 583
 FusionRed-ER fusion: 603-1397
 Start codon (ATG): 603-605
 Calreticulin signal sequence: 603-653
 FusionRed: 669-1364
 ER retention signal (KDEL): 1383-1394
 Stop codon: 1395-1397
 SV40 early mRNA polyadenylation signal
 Polyadenylation signals: 1589-1594 & 1618-1623
 mRNA 3' ends: 1627 & 1639
 f1 single-strand DNA origin: 1686-2141
 Bacterial promoter for expression of Kan^r gene
 -35 region: 2203-2208; -10 region: 2226-2231
 Transcription start point: 2238
 SV40 origin of replication: 2482-2617
 SV40 early promoter
 Enhancer (72-bp tandem repeats): 2315-2386 & 2387-2458
 21-bp repeats: 2462-2482, 2483-2503 & 2505-2525
 Early promoter element: 2538-2544
 Major transcription start points: 2534, 2572, 2578 & 2583
 Kanamycin/neomycin resistance gene
 Neomycin phosphotransferase coding sequences:
 Start codon (ATG): 2666-2668; Stop codon: 3458-3460
 G->A mutation to remove Pst I site: 2848
 C->A (Arg to Ser) mutation to remove BssH II site: 3194
 Herpes simplex virus (HSV) thymidine kinase (TK) polyadenylation signal
 Polyadenylation signals: 3696-3701 & 3709-3714
 pUC plasmid replication origin: 4045-4688

Product	Cat.#	Size
pFusionRed-ER vector	FP420	20 µg
Vector type	mammalian expression vector	
Reporter	FusionRed	
Reporter codon usage	mammalian	
Promoter for FusionRed	P _{CMV IE}	
Host cells	mammalian	
Selection	prokaryotic - kanamycin eukaryotic - neomycin (G418)	
Replication	prokaryotic - pUC ori eukaryotic - SV40 ori	
Use	red fluorescent labeling of the lumen of the endoplasmic reticulum	

Vector description

pFusionRed-ER is a mammalian expression vector intended for red fluorescent labeling of the lumen of the endoplasmic reticulum (ER) [Roderick et al. 1997]. The vector encodes red fluorescent protein FusionRed containing ER targeting signal (calreticulin signal sequence [Fliegel et al. 1989]) fused to the FusionRed N-terminus and ER retention signal (KDEL sequence [Munro and Pelham 1987]) fused to the FusionRed C-terminus.

FusionRed codon usage is optimized for high expression in mammalian cells (humanized) [Haas et al. 1996].

pFusionRed-ER vector can be used as a source of FusionRed-ER hybrid sequence. The vector backbone contains unique restriction sites that permit its excision and further insertion into expression vector of choice.

Note: The plasmid DNA was isolated from dam⁺-methylated *E. coli*. Therefore some restriction sites are blocked by methylation. If you wish to digest the vector using such sites you will need to transform the vector into a dam⁻ host and make fresh DNA.

The vector backbone contains immediate early promoter of cytomegalovirus (P_{CMV IE}) for protein expression, SV40 origin for replication in mammalian cells expressing SV40 T-antigen, pUC origin of replication for propagation in *E. coli*, and f1 origin for single-stranded DNA production. SV40 polyadenylation signals (SV40 poly A) direct proper processing of the 3'-end of the reporter mRNA.

SV40 early promoter (P_{SV40}) provides neomycin resistance gene (Neo^r) expression to select stably transfected eukaryotic cells using G418. Bacterial promoter (P) provides kanamycin resistance gene expression (Kan^r) in *E. coli*. Kan^r/Neo^r gene is linked with herpes simplex virus (HSV) thymidine kinase (TK) polyadenylation signals.

Expression in mammalian cells

pFusionRed-ER vector can be transfected into mammalian cells by any known transfection method. CMV promoter provides strong, constitutive expression of the FusionRed-ER fusion in eukaryotic cells. If required, stable transformants can be selected using G418 [Gorman 1985].

Propagation in *E. coli*

Suitable host strains for propagation in *E. coli* include DH5alpha, HB101, XL1-Blue, and other general purpose strains. Plasmid incompatibility group is pMB1/ColE1. The vector confers resistance to kanamycin (30 µg/ml) to *E. coli* hosts. Copy number in *E. coli* is about 500.

References

- Fliegel, L. et al. (1989) "Molecular cloning of the high affinity calcium-binding protein (calreticulin) of skeletal muscle sarcoplasmic reticulum." *J Biol Chem*, 264 (36): 21522-8 / pmid: 2600080
- Gorman, C. (1985). "High efficiency gene transfer into mammalian cells." In: *DNA cloning: A Practical Approach, Vol. II*. Ed. by Glover. (IRL Press, Oxford, U.K.) Pp. 143-190.
- Haas, J. et al. (1996) "Codon usage limitation in the expression of HIV-1 envelope glycoprotein." *Curr Biol*, 6 (3): 315-324 / pmid: 8805248
- Munro, S. and HR. Pelham (1987) "A C-terminal signal prevents secretion of luminal ER proteins." *Cell*, 48 (5): 899-907 / pmid: 3545499
- Roderick, HL. et al. (1997) "Nuclear localisation of calreticulin in vivo is enhanced by its interaction with glucocorticoid receptors." *FEBS Lett*, 405 (2): 181-185 / pmid: 9089287

Notice to Purchaser:

FusionRed-related materials (also referred to as "Products") are intended for research use only. The Products are covered by European Pat. 1994149 and other Evrogen Patents and/or Patent applications pending. By use of these Products, you accept the terms and conditions of the applicable Limited Use Label License #001: <http://www.evrogen.com/products/Evrogen-FP-license.shtml>.

The CMV promoter is covered under U.S. Patents 5,168,062 and 5,385,839, and its use is permitted for research purposes only. Any other use of the CMV promoter requires a license from the University of Iowa Research Foundation, 214 Technology Innovation Center, Iowa City, IA 52242.

MSDS information is available at <http://www.evrogen.com/MSDS.shtml>