



## peNirFP-N vector

The vector sequence has been compiled using the information from sequence databases, published literature, and other sources, together with partial sequences obtained by Evrogen. This vector has not been completely sequenced.



For vector sequence, please visit our Web site at <http://www.evrogen.com/products/vectors.shtml>

### Multiple cloning site (MCS)



### Location of features

$P_{CMVIE}$ : 1-589  
 Enhancer region: 59-465  
 TATA box: 554-560  
 Transcription start point: 583  
 MCS: 591-671  
 NirFP  
 Kozak consensus translation initiation site: 672-682  
 Start codon (ATG): 679-681; Stop codon: 1381-1383  
 Fragment of human beta globin (HBB) gene: 1392-2510  
 Last 35 bp of HBB exon 2: 1392-1426  
 HBB intron 2: 1427-2277  
 First 233 bp of HBB exon 3: 2278-2510  
 SV40 early mRNA polyadenylation signal  
 Polyadenylation signals: 2652-2657 & 2681-2686  
 mRNA 3' ends: 2690 & 2702  
 f1 single-strand DNA origin: 2749-3204  
 Bacterial promoter for expression of Kan<sup>r</sup> gene  
 -35 region: 3266-3271; -10 region: 3289-3294  
 Transcription start point: 3301  
 SV40 origin of replication: 3545-3680  
 SV40 early promoter  
 Enhancer (72-bp tandem repeats): 3378-3449 & 3450-3521  
 21-bp repeats: 3525-3545, 3546-3566 & 3568-3588  
 Early promoter element: 3601-3607  
 Major transcription start points: 3597, 3635, 3641 & 3646  
 Kanamycin/neomycin resistance gene  
 Neomycin phosphotransferase coding sequences:  
 Start codon (ATG): 3729-3731; Stop codon: 4521-4523  
 G->A mutation to remove Pst I site: 3911  
 C->A (Arg to Ser) mutation to remove BssH II site: 4257  
 Herpes simplex virus (HSV) thymidine kinase (TK) polyadenylation signal  
 Polyadenylation signals: 4759-4764 & 4772-4777  
 pUC plasmid replication origin: 5108-5751

### References

Gorman, C. (1985). "High efficiency gene transfer into mammalian cells." In: *DNA cloning: A Practical Approach, Vol. II*. Ed. by Glover. (IRL Press, Oxford, U.K.) Pp. 143-190.

Haas, J. et al. (1996) "Codon usage limitation in the expression of HIV-1 envelope glycoprotein." *Curr Biol*, 6 (3): 315-324 / pmid: 8805248

Kozak, M. (1987) "An analysis of 5'-noncoding sequences from 699 vertebrate messenger RNAs." *Nucleic Acids Res*, 15 (20): 8125-8148 / pmid: 3313277

### Notice to Purchaser:

NirFP-related materials (also referred to as "Products") are intended for research use only. The Products are covered by U.S. Pat. 7,972,834; 8,138,320; European Pat. 1994149; and other Evrogen Patents and/or Patent applications pending. By use of these Products, you accept the terms and conditions of the applicable Limited Use Label License #001: <http://www.evrogen.com/products/Evrogen-FP-license.shtml>.

The CMV promoter is covered under U.S. Patents 5,168,062 and 5,385,839, and its use is permitted for research purposes only. Any other use of the CMV promoter requires a license from the University of Iowa Research Foundation, 214 Technology Innovation Center, Iowa City, IA 52242.

MSDS information is available at <http://www.evrogen.com/MSDS.shtml>

Product	Cat.#	Size
peNirFP-N vector	FP743	20 µg
Vector type	mammalian expression vector	
Reporter	NirFP	
Reporter codon usage	mammalian	
Promoter for NirFP	$P_{CMVIE}$	
Host cells	mammalian	
Selection	prokaryotic - kanamycin eukaryotic - neomycin (G418)	
Replication	prokaryotic - pUC ori eukaryotic - SV40 ori	
Use	NirFP expression in mammalian cells; generation of fusions to the NirFP N-terminus	

### Vector description

peNirFP-N is a mammalian expression vector encoding near-infrared fluorescent protein NirFP. The vector allows generation of fusions to the NirFP N-terminus and expression of NirFP fusions or NirFP alone in eukaryotic (mammalian) cells.

NirFP codon usage is optimized for high expression in mammalian cells (humanized) [Haas et al. 1996]. To increase mRNA translation efficiency, Kozak consensus translation initiation site is generated upstream of the NirFP coding sequence [Kozak 1987]. Fragments of exons 2 and 3 and intron 2 of human beta globin gene are added in the 3' UTR of NirFP coding sequence in order to increase the protein expression level. Multiple cloning site (MCS) is located between  $P_{CMVIE}$  and NirFP coding sequence.

The vector backbone contains immediate early promoter of cytomegalovirus ( $P_{CMVIE}$ ) for protein expression, SV40 origin for replication in mammalian cells expressing SV40 T-antigen, pUC origin of replication for propagation in *E. coli*, and f1 origin for single-stranded DNA production. SV40 polyadenylation signals (SV40 poly A) direct proper processing of the 3'-end of the reporter mRNA.

SV40 early promoter ( $P_{SV40}$ ) provides neomycin resistance gene (Neo<sup>r</sup>) expression to select stably transfected eukaryotic cells using G418. Bacterial promoter (P) provides kanamycin resistance gene expression (Kan<sup>r</sup>) in *E. coli*. Kan<sup>r</sup>/Neo<sup>r</sup> gene is linked with herpes simplex virus (HSV) thymidine kinase (TK) polyadenylation signals.

### Generation of NirFP fusion proteins

A localization signal or a gene of interest can be cloned into MCS of the vector. It will be expressed as a fusion to the NirFP N-terminus when inserted in the same reading frame as NirFP and no in-frame stop codons are present. The inserted sequence should contain an initiating ATG codon. NirFP-tagged fusions retain fluorescent properties of the native protein allowing fusion localization *in vivo*. Unmodified vector will express NirFP when transfected into eukaryotic (mammalian) cells.

**Note:** The plasmid DNA was isolated from dam<sup>+</sup>-methylated *E. coli*. Therefore some restriction sites are blocked by methylation. If you wish to digest the vector using such sites you will need to transform the vector into a dam<sup>-</sup> host and make fresh DNA.

### Expression in mammalian cells

peNirFP-N vector can be transfected into mammalian cells by any known transfection method. CMV promoter provides strong, constitutive expression of NirFP or its fusions in eukaryotic cells. If required, stable transformants can be selected using G418 [Gorman 1985].

### Propagation in *E. coli*

Suitable host strains for propagation in *E. coli* include DH5alpha, HB101, XL1-Blue, and other general purpose strains. Plasmid incompatibility group is pMB1/ColE1. The vector confers resistance to kanamycin (30 µg/ml) to *E. coli* hosts. Copy number in *E. coli* is about 500.