



pPA-TagRFP-N vector

The vector sequence has been compiled using the information from sequence databases, published literature, and other sources, together with partial sequences obtained by Evrogen. This vector has not been completely sequenced.



For vector sequence, please visit our Web site at <http://www.evrogen.com/products/vectors.shtml>

Multiple cloning site (MCS)



Location of features

P_{CMV IE}: 1-589
 Enhancer region: 59-465
 TATA box: 554-560
 Transcription start point: 583
 MCS: 591-671
 PA-TagRFP
 Kozak consensus translation initiation site: 672-682
 Start codon (ATG): 679-681; Stop codon: 1378-1380
 SV40 early mRNA polyadenylation signal
 Polyadenylation signals: 1534-1539 & 1563-1568
 mRNA 3' ends: 1572 & 1584
 f1 single-strand DNA origin: 1631-2086
 Bacterial promoter for expression of Kan^r gene
 -35 region: 2148-2153; -10 region: 2171-2176
 Transcription start point: 2183
 SV40 origin of replication: 2427-2562
 SV40 early promoter
 Enhancer (72-bp tandem repeats): 2260-2331 & 2332-2403
 21-bp repeats: 2407-2427, 2428-2448 & 2450-2470
 Early promoter element: 2483-2489
 Major transcription start points: 2479, 2517, 2523 & 2528
 Kanamycin/neomycin resistance gene
 Neomycin phosphotransferase coding sequences:
 Start codon (ATG): 2611-2613; Stop codon: 3403-3405
 G->A mutation to remove Pst I site: 2793
 C->A (Arg to Ser) mutation to remove BssH II site: 3139
 Herpes simplex virus (HSV) thymidine kinase (TK) polyadenylation signal
 Polyadenylation signals: 3641-3646 & 3654-3659
 pUC plasmid replication origin: 3990-4633

References

Gorman, C. (1985). "High efficiency gene transfer into mammalian cells." In: *DNA cloning: A Practical Approach, Vol. II*. Ed. by Glover. (IRL Press, Oxford, U.K.) Pp. 143-190.

Haas, J. et al. (1996) "Codon usage limitation in the expression of HIV-1 envelope glycoprotein." *Curr Biol*, 6 (3): 315-324 / pmid: 8805248

Kozak, M. (1987) "An analysis of 5'-noncoding sequences from 699 vertebrate messenger RNAs." *Nucleic Acids Res*, 15 (20): 8125-8148 / pmid: 3313277

Subach, FV et al. (2010) "Bright monomeric photoactivatable red fluorescent protein for two-color super-resolution sptPALM of live cells." *J Am Chem Soc*, 132 (18): 6481-91 / pmid: 20394363

Notice to Purchaser:

PA-TagRFP-related materials (also referred to as "Products") are intended for research use only. The Products are covered by European Pat. 1994149 and other Evrogen Patents and/or Patent applications pending. By use of these Products, you accept the terms and conditions of the applicable Limited Use Label License #001: <http://www.evrogen.com/products/Evrogen-FP-license.shtml>.

The CMV promoter is covered under U.S. Patents 5,168,062 and 5,385,839, and its use is permitted for research purposes only. Any other use of the CMV promoter requires a license from the University of Iowa Research Foundation, 214 Technology Innovation Center, Iowa City, IA 52242.

MSDS information is available at <http://www.evrogen.com/MSDS.shtml>

Product	Cat.#	Size
pPA-TagRFP-N vector	FP812	20 µg
Vector type	mammalian expression vector	
Reporter	PA-TagRFP	
Reporter codon usage	mammalian	
Promoter for PA-TagRFP	P _{CMV IE}	
Host cells	mammalian	
Selection	prokaryotic - kanamycin eukaryotic - neomycin (G418)	
Replication	prokaryotic - pUC ori eukaryotic - SV40 ori	
Use	PA-TagRFP expression in mammalian cells; generation of fusions to the PA-TagRFP N-terminus	

Vector description

pPA-TagRFP-N is a mammalian expression vector encoding photoactivatable red fluorescent protein PA-TagRFP. The vector allows generation of fusions to the PA-TagRFP N-terminus and expression of PA-TagRFP fusions or PA-TagRFP alone in eukaryotic (mammalian) cells.

Note: The pPA-TagRFP-N vector encodes the PA-TagRFP protein in which the amino acid sequence at N-terminus is modified compared to the originally reported sequence [Subach et al. 2010]. These modifications do not influence the fluorescent properties of the PA-TagRFP, but improves its performance in fusions.

PA-TagRFP codon usage is optimized for high expression in mammalian cells (humanized) [Haas et al. 1996]. To increase mRNA translation efficiency, Kozak consensus translation initiation site is generated upstream of the PA-TagRFP coding sequence [Kozak 1987]. Multiple cloning site (MCS) is located between P_{CMV IE} and PA-TagRFP coding sequence.

The vector backbone contains immediate early promoter of cytomegalovirus (P_{CMV IE}) for protein expression, SV40 origin for replication in mammalian cells expressing SV40 T-antigen, pUC origin of replication for propagation in *E. coli*, and f1 origin for single-stranded DNA production. SV40 polyadenylation signals (SV40 poly A) direct proper processing of the 3'-end of the reporter mRNA.

SV40 early promoter (P_{SV40}) provides neomycin resistance gene (Neo^r) expression to select stably transfected eukaryotic cells using G418. Bacterial promoter (P) provides kanamycin resistance gene expression (Kan^r) in *E. coli*. Kan^r/Neo^r gene is linked with herpes simplex virus (HSV) thymidine kinase (TK) polyadenylation signals.

Generation of PA-TagRFP fusion proteins

A localization signal or a gene of interest can be cloned into MCS of the vector. It will be expressed as a fusion to the PA-TagRFP N-terminus when inserted in the same reading frame as PA-TagRFP and no in-frame stop codons are present. The inserted sequence should contain an initiating ATG codon. PA-TagRFP-tagged fusions retain fluorescent properties of the native protein allowing fusion localization *in vivo*. Unmodified vector will express PA-TagRFP when transfected into eukaryotic (mammalian) cells.

Note: The plasmid DNA was isolated from dam⁺-methylated *E. coli*. Therefore some restriction sites are blocked by methylation. If you wish to digest the vector using such sites you will need to transform the vector into a dam⁻ host and make fresh DNA.

Expression in mammalian cells

pPA-TagRFP-N vector can be transfected into mammalian cells by any known transfection method. CMV promoter provides strong, constitutive expression of PA-TagRFP or its fusions in eukaryotic cells. If required, stable transformants can be selected using G418 [Gorman 1985].

Propagation in *E. coli*

Suitable host strains for propagation in *E. coli* include DH5alpha, HB101, XL1-Blue, and other general purpose strains. Plasmid incompatibility group is pMB1/ColE1. The vector confers resistance to kanamycin (30 µg/ml) to *E. coli* hosts. Copy number in *E. coli* is about 500.