


## pTurboRFP-N vector

The vector sequence has been compiled using the information from sequence databases, published literature, and other sources, together with partial sequences obtained by Evrogen. This vector has not been completely sequenced.



For vector sequence, please visit our Web site at:  
<http://www.evrogen.com/support/vector-info.shtml>

### Multiple cloning site (MCS)

<i>Afe I</i>	<i>Xba I</i>	<i>Hind III</i>	<i>Pst I</i>	<i>Kpn I</i>	<i>Apa I</i>	<i>BamH I</i>	<i>TurboRFP</i>	
... G.CTA.GCG.CTA.CCG.GAC.TCA.GAT.CTC.GAG.CTC.AAG.CTT.CGA.ATT.CTG.CAG.TCG.ACG.GTA.CCG.CGG.GCC.GCG.GAT.CCA.CCG.GTC.GCC.ACC.ATG.A ...	Nhe I	Bgl II	Sac I	EcoR I	Sai I	Sac II	Sma I/Xma I	Age I

### Location of features

P<sub>CMV IE</sub>: 1-589  
 Enhancer region: 59-465  
 TATA box: 554-560  
 Transcription start point: 583  
 MCS: 591-671  
 Turbo-RFP  
 Kozak consensus translation initiation site: 672-682  
 Start codon (ATG): 679-681; Stop codon: 1375-1377  
 SV40 early mRNA polyadenylation signal  
 Polyadenylation signals: 1528-1533 & 1557-1562  
 mRNA 3' ends: 1566 & 1578  
 f1 single-strand DNA origin: 1625-2080  
 Eukaryotic promoter for expression of Kan' gene  
 -35 region: 2142-2147; -10 region: 2165-2170  
 Transcription start point: 2177  
 SV40 origin of replication: 2421-2556  
 SV40 early promoter  
 Enhancer (72-bp tandem repeats): 2254-2325 & 2326-2397  
 21-bp repeats: 2401-2421, 2422-2442 & 2444-2464  
 Early promoter element: 2477-2483  
 Major transcription start points: 2473, 2511, 2517 & 2522  
 Kanamycin/neomycin resistance gene  
 Neomycin phosphotransferase coding sequences:  
 Start codon (ATG): 2605-2607; Stop codon: 3397-3399  
 G->A mutation to remove Pst I site: 2787  
 C->A (Arg to Ser) mutation to remove BssH II site: 3133  
 Herpes simplex virus (HSV) thymidine kinase (TK) polyadenylation signal  
 Polyadenylation signals: 3635-3640 & 3648-3653  
 pUC plasmid replication origin: 3984-4627

### References

- Gorman (1985). "High efficiency gene transfer into mammalian cells." In: *DNA cloning: A Practical Approach, Vol. II*. Ed. by Glover. (IRL Press, Oxford, U.K.) Pp. 143-190.  
 Haas et al. (1996) "Codon usage limitation in the expression of HIV-1 envelope glycoprotein." *Curr Biol*, 6 (3): 315-324 / pmid: 8805248  
 Kozak (1987) "An analysis of 5'-noncoding sequences from 699 vertebrate messenger RNAs." *Nucleic Acids Res*, 15 (20): 8125-8148 / pmid: 3313277

Product	Cat.#	Size
pTurboRFP-N vector	FP232	20 µg
The price does not include delivery. The price varies in different countries. Please contact your local distributor for exact prices and delivery information.		
Vector type	mammalian expression vector	
Reporter	TurboRFP	
Reporter codon usage	mammalian	
Promoter for TurboRFP	P <sub>CMV IE</sub>	
Host cells	mammalian	
Selection	prokaryotic - kanamycin eukaryotic - neomycin (G418)	
Replication	prokaryotic - pUC ori eukaryotic - SV40 ori	
Use	TurboRFP expression in mammalian cells; generation of fusions to the TurboRFP N-terminus	

### Vector description

TurboRFP-N is a mammalian expression vector encoding red (orange) fluorescent protein TurboRFP. The vector allows generation of fusions to the TurboRFP N-terminus and expression of TurboRFP fusions or TurboRFP alone in eukaryotic (mammalian) cells.

TurboRFP codon usage is optimized for high expression in mammalian cells (humanized) [Haas et al. 1996]. To increase mRNA translation efficiency, Kozak consensus translation initiation site is generated upstream of TurboRFP sequence [Kozak 1987]. Multiple cloning site (MCS) is located between P<sub>CMV IE</sub> and TurboRFP coding sequence.

The vector backbone contains immediate early promoter of cytomegalovirus (P<sub>CMV IE</sub>) for protein expression, SV40 origin for replication in mammalian cells expressing SV40 T-antigen, pUC origin of replication for propagation in *E. coli* and f1 origin for single-stranded DNA production. SV40 polyadenylation signals (SV40 poly A) direct proper processing of the 3'-end of the reporter mRNA.

SV40 early promoter (P<sub>SV40</sub>) provides neomycin resistance gene (Neo<sup>r</sup>) expression to select stably transfected eukaryotic cells using G418. Bacterial promoter (P) provides kanamycin resistance gene expression (Kan<sup>r</sup>) in *E. coli*. Kan<sup>r</sup>/Neo<sup>r</sup> gene is linked with herpes simplex virus (HSV) thymidine kinase (TK) polyadenylation signals.

### Generation of TurboRFP-tagged fusions

A localization signal or a gene of interest should be cloned into MCS of the vector. It will be expressed as a fusion to the TurboRFP N-terminus when inserted in the same reading frame as TurboRFP and no in-frame stop codons are present. The inserted sequence should contain an initiating ATG codon. TurboRFP-tagged fusions retain fluorescent properties of the native protein allowing fusion localization *in vivo*. Unmodified vector will express TurboRFP, when transfected into eukaryotic (mammalian) cells.

**Note:** The plasmid DNA was isolated from dam<sup>+</sup>-methylated *E. coli*. Therefore some restriction sites are blocked by methylation. If you wish to digest the vector using such sites you will need to transform the vector into a dam<sup>-</sup> host and make fresh DNA.

Despite its dimeric structure, TurboRFP is still suitable for generation of fusions with proteins of interest, however we recommend to use TagFPs for these purposes.

### Expression in mammalian cells

pTurboRFP-N vector can be transfected into mammalian cells by any known transfection method. If required, stable transformants can be selected using G418 [Gorman 1985].

### Propagation in *E. coli*

Suitable host strains for propagation in *E. coli* include DH5alpha, HB101, XL1-Blue, and other general purpose strains. Plasmid incompatibility group is pMB1/ColE1. The vector confers resistance to kanamycin (30 µg/ml) to *E. coli* hosts. Copy number in *E. coli* is about 500.

### Notice to Purchaser:

Evrogen Fluorescent Protein Products (the Products) are intended for research use only. The Products are covered by U.S. Pat. 7,417,131 and other Evrogen Patents and/or Patent applications pending. By use of these Products, you accept the terms and conditions of the applicable Limited Use Label License.

The CMV promoter is covered under U.S. Patents 5,168,062 and 5,385,839, and its use is permitted for research purposes only. Any other use of the CMV promoter requires a license from the University of Iowa Research Foundation, 214 Technology Innovation Center, Iowa City, IA 52242.

**MATERIAL SAFETY DATA SHEET INFORMATION:** To the best of our knowledge, these products do not require a Material Safety Data Sheet. However, all the properties of these products (and, if applicable, each of their components) have not been thoroughly investigated. Therefore, we recommend that you use gloves and eye protection, and wear a laboratory coat when working with these products.